

LULY

LULICONAZOLE INN

Compositions:

Each gram cream contains Luliconazole INN 10 mg.

Pharmacology:

Luliconazole is a broad-spectrum azole antifungal that acts by inhibiting the enzyme lanosterol demethylase and blocks ergosterol synthesis which is an important component of fungal cell membrane. Thus, it exerts strong fungicidal activity both in-vitro and in-vivo against dermatophytes, *Candida albicans*, *Malassezia* spp., *Trichophyton* & *Epidermophyton* spp.

Dosage And Administration:

Interdigital Tinea Pedis: Luly Cream should be applied to the affected and immediate surrounding area once a day for two weeks. Tinea Cruris and Tinea Corporis: Luly Cream should be applied to the affected skin and immediate surrounding area once a day for one week.

Contraindications:

None.

Warning And Precaution:

Luliconazole is recommended for topical use only. It is not intended for ophthalmic, oral or intravaginal use.

Side Effects:

Contact dermatitis and cellulitis may occur. Application site reactions were observed in less than 1% of subjects in clinical trial.

Use in Pregnancy and Lactation:

Pregnancy: Pregnancy category C. There are no adequate and well controlled studies in pregnant women, Luliconazole should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Lactation: It is not known whether Luliconazole is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when Luliconazole cream is applied to women who are breastfeeding.

Drug Interaction:

No information was found.

Overdosage:

No information was found.

Storage:

Store below 30° C. Keep away from light and out of the reach of children.

Packing:

Each pack has a laminated tube containing 20 gm cream.

Manufactured By:

The IBN SINA Pharmaceutical Industry PLC.

Shafipur, Gazipur, Bangladesh.