

Ribosina

Riboflavin

Composition:

Each tablet contains Riboflavin BP 5 mg.

Indication:

To prevent and to treat deficiencies of vitamin B2 such as cheilosis, angular stomatitis, glossitis, keratitis, seborrhoeic dermatitis.

Dosage and Administration:

- For treating low levels of riboflavin (riboflavin deficiency) in adults: 5-30 mg of riboflavin daily in divided doses.
- For preventing migraine headaches: 400 mg of riboflavin (Vitamin B2) per day. It may take up to three months to get best results.
- For preventing cataracts: approximately 2.6 mg of riboflavin (Vitamin B2)

Contraindication:

Hypersensitivity to Riboflavin.

Precaution:

Riboflavin is non-toxic. No warnings or *precautions* have been established .

Side Effect:

Riboflavin is **LIKELY SAFE** for most people. In some people, Riboflavin can cause the urine to turn a yellow-orange color. When taken in high doses, Riboflavin might cause diarrhea, an increase in urine, and other side effects.

Drug Interaction:

Drying medications (Anticholinergic drugs), Medications for depression (Tricyclic antidepressants), Phenobarbital, Probenecid.

Use in Pregnancy and Lactation:

Riboflavin is **LIKELY SAFE** for pregnant or breast-feeding women when taken in the amounts recommended.

Storage Condition:

Store at a cool and dry place. Protect from light.

Presentation:

Each box contains 50×10's tablet in Alu-PVC strip.

Manufactured by:



The IBN SINA Pharmaceutical Industry Ltd.
Shafipur, Gazipur, Bangladesh.