

Veesina

Vitamin-C
Chewable Tablet

Composition

Each **Veesina** tablet contains Ascorbic Acid BP 125 mg.
Sodium Ascorbate USP 140.62 mg (equivalent to 125 mg of Ascorbic acid).

Description:

Vitamin-C (**Veesina**) is an important constituent of the body for biological oxidation-reduction process. It is essential in the synthesis of collagen and intracellular materials. Besides it has an essential role in the formation of soft bones, maturation of teeth by formation of dentin, conversion of folic acid to folinic acid and for the remedy of injury and wounds. Also it plays an important role in haemoglobin synthesis and in maturation of RBC. Recent studies indicated that the old men who have low ascorbic acid level in blood have higher risk of stroke. So, long term intake of ascorbic acid at a level higher than normal results in a greater decrease in risk of stroke.

Indication

- 1 Vitamin-C deficiency diseases, such as: scurvy, dental caries, pyorrhea, haemorrhage, dental haemorrhage, anaemia etc.
- 2 To increase body resistance.
- 3 To increase iron absorption from iron containing foods.
- 4 During pregnancy and for women taking contraceptive pill.
- 5 For old-ster and for chain smoker.
- 6 For normal wound healing and post-operative wound healing, as it helps in tissue repairing and in formation of collagen.

Dosage and Administration

Scurvy: 1-4 tablets daily or as advised by the physician.

Wound healing : 1-2 tablets daily for 7-10 days.

Ulcer due to severe burn: 4-8 tablets daily.

Other cases: 1-2 tablets daily.

Precaution

During pregnancy and lactation, Vitamin-C should be given with caution & must not be given in higher dose.

Side effect

Higher dose may cause diarrhoea, headache, nausea. Stones of oxalate and urate may form in kidney, if urine becomes acidic.

Drug interaction

Bleomycin, Erythromycin, Chloramphenicol, Chlorthiazide Sodium, Hydrocortison Sodium etc.

How supplied

Veesina Tablet : Box contains 10 x 10's tablets in aluminium strip.