

MYOSON

TOLPERISONE INN

Compositions:

Myoson 50 Tablet: Each film-coated tablet contains Tolperisone Hydrochloride INN 50 mg.
Myoson 100 Tablet: Each film-coated tablet contains Tolperisone Hydrochloride INN 100 mg.

Pharmacology:

Tolperisone Hydrochloride is a centrally acting muscle relaxant. It effectively reduces muscular hypertonicity, improves peripheral blood flow by vasodilatation and reduces pain.

Dosage And Administration:

The recommended dose is as followings: Adult: 50-150 mg three times daily according to the individual requirement and tolerance of the patient. Children (from 3 months to 6 years): 5-10 mg/kg/day in three divided doses. Children (from 6 to 14 years): 2-4 mg/kg/day in three divided doses. One should follow doctor's instructions regarding the dosage of Myoson.

Contraindications:

Hypersensitivity to any component of the product • Myasthenia gravis • Nursing mother

Warning And Precaution:

In case of children, the prescribed dose and duration of treatment should closely be observed.

Side Effects:

It may cause excessive urticaria, erythema. Also may lead to GI upset with abdominal pain, nausea, vomiting, diarrhea, flatulence or dryness of mouth.

Use in Pregnancy and Lactation:

Tolperisone Hydrochloride should be used in pregnancy according to physician's advice. Tolperisone Hydrochloride should not be used during breast-feeding.

Drug Interaction:

Concomitant use of methocarbamol with Tolperisone hydrochloride has been reported to cause disturbance of visual accommodation.

Overdosage:

In the event of over dosage supportive therapy may be substituted.

Storage:

Keep in a cool (below 30°C) and dry place, protected from light. Keep out of the reach of children.

Packing:

Myoson-50: Each box contains 10 x 10 tablets in Alu-PVC blister strips. Myoson-100: Each box contains 5 x 10 tablets in Alu-PVC blister strips.

Manufactured By:

The IBN SINA Pharmaceutical Industry PLC.
Shafipur, Gazipur, Bangladesh.