

# NEXYL

TREANEXAMIC ACID BP

## **Compositions:**

Nexyl Capsule : Each capsule contains Tranexamic Acid BP 500 mg.

## **Pharmacology:**

Tranexamic acid has a strong inhibitory effect on the activation of plasminogen, i.e. the conversion of plasminogen to plasmin, in the fibrinolytic system. The half-life is 1-2 hours. Plasma protein binding is 3% at therapeutic levels. The plasma protein binding seems fully accounted by its binding to plasminogen. Tranexamic acid is excreted unchanged in the urine. Tranexamic acid is rapidly absorbed from the gastrointestinal tract. Maximum serum levels are reached within 2-3 hours. After oral administration, about 40% of the dose is excreted in the urine during the first 24 hours. After intravenous administration, 45% of the dose is excreted in the urine during the first day.

## **Dosage And Administration:**

2-3 capsules 3 times daily for maximum 5 days during menstruation.

## **Contraindications:**

Active thromboembolic disease, such as deep vein thrombosis, pulmonary embolism and cerebral thrombosis Subarachnoid haemorrhage Hypersensitivity to tranexamic acid or any of the ingredients.

## **Warning And Precaution:**

Patients with irregular menstrual bleeding, patients with a high risk of thrombosis (a previous thromboembolic event and a family history of thromboembolic disease) should use it only if there is a strong medical indication and under strict medical supervision. Patients with disseminated intravascular coagulation (DIC), who require treatment with it must be under the strict supervision of a physician experienced in treating this disorder. In the long-term treatment of patients, regular eye examination should be performed. If a colour vision disorder occurs during the course of treatment, the drug should be discontinued.

## **Side Effects:**

Dose-dependent gastrointestinal discomfort is the most commonly reported undesirable effect, but it is usually of mild and temporary in nature. Allergic skin reactions have been reported as an uncommon undesirable effect. Hypotension may occur after fast injection.

## **Use in Pregnancy and Lactation:**

Pregnancy: Tranexamic acid crosses the placenta. Clinical experience of use in pregnant women is limited. Animal studies have not supplied any evidence of an increased incidence of foetal damage. Lactation: Tranexamic acid is excreted into breast milk, but it is not likely to influence the child at therapeutic doses.

## **Drug Interaction:**

Tranexamic Acid is a synthetic Amino Acid that is incompatible with solutions containing penicillins (eg: Benzyl penicillin). Thrombolytic drugs like Streptokinase & Urokinase antagonise the antifibrinolytic action of Tranexamic Acid. The potential for thrombus formation may be increased by concomitant administration of estrogen containing drugs, like oral contraceptives. Direct admixture of Tranexamic Acid with whole blood should be avoided during Transfusion.

**Overdosage:**

No data available

**Storage:**

Stored between 15°C to 30° C and protected from light and moisture.

**Packing:**

Nexyl Capsule: Each box contains 62 x 10's capsules in blister pack.

Manufactured By:

The IBN SINA Pharmaceutical Industry PLC.

Shafipur, Gazipur, Bangladesh.