

## UNISALINE

SODIUM CHLORIDE; POTASSIUM CHLORIDE; TRISODIUM CITRATE DIHYDRATE &  
GLUCOSE ANHYDROUS

### **Compositions:**

Each sachet contains Sodium Chloride- 1.30 gm, Potassium Chloride - 0.75 gm, Sodium Citrate(As dehydrate)-1.45 gm, Anhydrous glucose – 6.75 gm,

### **Pharmacology:**

Oral Rehydration Saline. Oral rehydration therapy is a treatment for the symptoms of dehydration.

### **Dosage And Administration:**

The amount of water removed from the body, equivalent quantity of mixture should be given. After each loose motion... Children below 2 years : 50-100 ml (10-20 teaspoonful) of prepared Unisaliner. Children between 2-10 years : 100-200 ml (20-40 teaspoonful) prepared Unisaliner. Children above 10 years and adults : 200-400 ml (1-2 glass) prepared Unisaliner.

### **Warning And Precaution:**

ORS should not be mixed with hot water and the mixture should not be heated. The mixture must be discarded after 12 hours from preparation.

### **Side Effects:**

No data is available

### **Drug Interaction:**

No data is available

### **Overdosage:**

No data is available

### **Storage:**

Store in a cool and dry place, away from light. Keep out of the reach of children.

### **Packing:**

Each box contains 20 sachets.

Manufactured By:

The IBN SINA Pharmaceutical Industry PLC.  
Shafipur, Gazipur, Bangladesh.