

BROMOFEN

BROMFENAC INN

Compositions:

Bromofen eye drops : Each ml contains Bromfenac sodium INN equivalent to Bromfenac 0.9 mg.

Pharmacology:

Bromofen eye drops is a sterile, topical, non steroidal anti-inflammatory drug (NSAID) for ophthalmic use.

Dosage And Administration:

For the treatment of post-operative inflammation in patients who have undergone cataract extraction, one drop of Bromfenac ophthalmic solution should be applied to the affected eye(s) two times daily beginning 24 hours after cataract surgery and continuing through the first 2 weeks of the post-operative period.

Contraindications:

Bromfenac ophthalmic solution is contraindicated in patients with known hypersensitivity to any ingredient in the formulation.

Warning And Precaution:

It is recommended that Bromfenac ophthalmic solution be used with caution in patients with known bleeding tendencies or who are receiving other medications which may prolong bleeding time.

Side Effects:

The most commonly reported adverse experiences reported following use of bromfenac after cataract surgery include: abnormal sensation in eye, conjunctival hyperemia, eye irritation (including burning/stinging), eye pain, eye pruritus, eye redness, headache, and iritis. These events were reported in 2-7% of patients.

Use in Pregnancy and Lactation:

Pregnancy Category C. This drug should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Caution should be exercised when Bromfenac ophthalmic solution is administered to a nursing mother

Drug Interaction:

No information provided.

Overdosage:

No information provided.

Storage:

Store in a cool, dry place and protected from light. Keep out of reach of children. Discard the container 4 weeks after opening.

Packing:

Bromofen eye drops : 5 ml sterile solution in plastic dropper bottle.

Manufactured By:
The IBN SINA Pharmaceutical Industry PLC.
Shafipur, Gazipur, Bangladesh.