

EYLON

FLUROMETHOLON USP + TETRAZOLIN HYDROCHLORIDE USP

Compositions:

Eylon Eye Drops : Each ml contains Fluorometholone USP 1mg & Tetrazoline Hydrochloride USP 0.25 mg.

Pharmacology:

Fluorometholone, like all other antiinflammatory glucocorticoids, gives its antiinflammatory activity by inhibiting phospholypase A2, the first step in prostaglandin synthesis. It also inhibits the chemotactic infiltration of neutrophils at the site of inflammation. The major benefit of fluorometholone over other corticosteroids is that it is not only 40 times stronger antiinflammatory drug but also has no significant action in increasing intraocular pressure. Fluorometholone is more rapidly degraded in the tissue. Therefore, immuno suppressive action of Flurometholone is very less. Tetrazoline, an a-sympathomimetic, is a local vasoconstrictor which acts promptly and alleviates hyperemia, redness, swelling and irritation of conjunctiva.

Dosage And Administration:

Adults: Instill 1 drop 2-3 times daily into the conjunctival sac for first one to two days. The dose may be increased for severe and adult patient to one drop hourly. Children (above two years): Specific studies for children have not been performed yet. So, Eylon should be used with caution in children.

Contraindications:

Hypersensitivity to any component of this product, infectious conjunctivitis and keratitis, ulcerous condition of cornea, infection caused by virus, bacteria & fungus, glaucoma, stromal damage of the cornea or sclera, dry eyes specially keratoconjunctivitis sicca and children under two years.

Warning And Precaution:

This preparation is not for prolonged use. If long term treatment is necessary (more than 2 or 3 days) patients should be regularly examined with regard to intraocular pressure, systemic adverse reactions and secondary infections. The preparation should be used after a careful risk/benefit evaluation in case of patient with,-Severe blood circulation disease. - Metabolic disturbance. -Under treatment with monoamine oxidase inhibitors and other potentially hypertensive drugs.

Side Effects:

Slight burning sensation for a short time after application of this eye drops have been reported by a small number of patients. Tetrazoline may cause slight irritation, in some cases mydriasis, dry mucous membrane and in rare case due to vasoconstriction, palpitation, arrythmia, angina have been reported.

Use in Pregnancy and Lactation:

No controlled human studies are available, animal studies with corticosteroids have revealed undesirable effect on the fetus. So, using of this medicine during pregnancy & lactation is therefore not recommended.

Drug Interaction:

Eylon should not be administered concomitantly with monoamine oxidase inhibitors or tricyclic antidepressants which could increase blood pressure due to vasoconstricting effect.

Overdosage:

When the product is used as directed, there is almost no likelihood of an overdose. No information on overdose with fluorometholone is available. Overdosage with fluorometholone is unlikely to cause acute problems. The symptoms of acute overdose with tetrazoline are CNS, cardiac and psychiatric disturbances, mydriasis, cyanosis and fever. CNS functions may be inhibited under certain circumstances. The following measures are possible in case of accidental ingestion and the occurrence of symptoms of intoxication: administration of activated charcoal, gastric lavage, artificial ventilation with oxygen, use of phentolamine to lower blood pressure (5 mg in saline solution, given i.v.). Vasopressors are contraindicated. Antipyretic and anticonvulsive therapy can be administered as necessary.

Storage:

Should not be used after the date marked "EXP" on the pack. This must be kept out of the reach and sight of children.

Packing:

Eylon Eye Drops: Each dropper bottle contains 5ml of sterile eye drops.

Manufactured By:

The IBN SINA Pharmaceutical Industry PLC.

Shafipur, Gazipur, Bangladesh.