

NITASOL

NITAZOXANIDE INN

Compositions:

Nitasol Tablet: Each film coated tablet contains Nitazoxanide INN 500 mg. Nitasol 30 ml suspension: Each 5 ml reconstituted suspension contains Nitazoxanide INN 100 mg. Nitasol 60 ml suspension: Each 5 ml reconstituted suspension contains Nitazoxanide INN 100 mg.

Pharmacology:

Nitasol (Nitazoxanide), a synthetic antiprotozoal drug. It is well absorbed from gastrointestinal tract. It interferes with the Pyruvate Ferredoxin Oxidoreductase (PFOR) enzyme dependent electron transfer reaction which is essential to anaerobic glucose energy metabolism. This results in cell swelling, membrane damage resulting in dysfunction of the parasite.

Dosage And Administration:

Age 1-3 years: 5 ml (100 mg) twice daily for 3 days. Age 4-11 years: 10 ml (200 mg) twice daily for 3 days. Age >12 years: 25 ml or 1 tablet (500 mg) twice daily for 3 days. The suspension or tablet should be taken with food.

Contraindications:

Known hypersensitivity to Nitazoxanide or any other ingredient in the formulations.

Warning And Precaution:

Nitazoxanide should be administered with caution to patients with hepatic, renal and biliary disease.

Side Effects:

Nitazoxanide is generally well tolerated. However abdominal pain, diarrhea vomiting and headache have been reported rarely.

Use in Pregnancy and Lactation:

Pregnancy category B: This drug should be used during pregnancy only if clearly needed. Nursing mother: It is not known whether Nitazoxanide is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when Nitazoxanide is administered to a nursing woman.

Drug Interaction:

Nitazoxanide is highly bound to plasma protein (>99.9%). Therefore, caution should be used when administering Nitazoxanide concurrently with other highly plasma-protein bound drugs.

Overdosage:

Information on Nitazoxanide over dosage is not available. Single oral doses up to 4000 mg Nitazoxanide have been administered to healthy adults without significant adverse effects.

Storage:

Store in a cool and dry place & away from light and moisture. The reconstituted suspension should be used within 7 days of preparation.

Packing:

Nitasol Tablet: Box contains 12 tablets in Alu-Alu blister. Nitasol 30 ml suspension: Bottle containing dry powder to reconstitute 30 ml suspension. Nitasol 60 ml suspension: Bottle containing dry powder to reconstitute 60 ml suspension.

Manufactured By:
The IBN SINA Pharmaceutical Industry PLC.
Shafipur, Gazipur, Bangladesh.