

OSTORIN

GLUCOSAMINE SULFATE USP & DIACEREIN INN

Compositions:

Ostorin Tablet: Each tablet contains Glucosamine Sulfate 750 mg as Glucosamine Sulfate Potassium Chloride USP and Diacerein INN 50 mg.

Pharmacology:

Ostorin is the combination of 750 mg of Glucosamine sulfate and 50 mg Diacerein. Two symptomatic slow acting drug in Osteoarthritis. Diacerein is a new anti - inflammatory analgesic and antipyretic drug which directly inhibits IL - 1 beta synthesis and release in vitro and down modulates IL - 1 induced activities and have been shown to possess disease modifying effect in human subjects with finger joint and knee Osteoarthritis. Glucosamine Sulfate is a naturally occurring chemical found in the human body and it is commonly used in osteoarthritis.

Dosage And Administration:

Adult and Children 12 years and older: The usual dose of Ostorin is 1 tablet twice daily (12 hourly) taken with food.

Contraindications:

There are no known contraindications for Glucosamine. But proven hypersensitivity to Glucosamine is a contraindication. Diacerein is contraindicated in pregnancy, lactation and hypersensitivity to anthraquinone derivatives.

Warning And Precaution:

Caution should be practiced when administering Ostorin in those patients who are allergic to Glucosamine and/or Diacerein or to shellfish (shrimp, crab); who have diabetes, asthma, alcohol dependence or liver disease. Before taking Ostorin complete blood count, liver function and urinalysis should be monitored regularly.

Side Effects:

Diarrhea, constipation, epigastric pain, heart burn, nausea, vomiting, headache, skin rashes, drowsiness, intense yellow coloring of urine may be associated with the active components of Ostorin.

Use in Pregnancy and Lactation:

Glucosamine and Diacerein are contraindicated during pregnancy and breastfeeding.

Drug Interaction:

Absorption of Diacerein is reduced if taken with aluminium and/or magnesium hydroxide antacid. Diacerein also increases risk of diarrhea if taken with laxatives, antibiotics. Avoid co-administration with fibers and phytic acids. Diacerein has shown hypersensitivity to anthraquinone derivatives.

Overdosage:

There is no data available regarding overdose of this combination.

Storage:

Store at a cool and dry place, away from light. Keep out of reach of children.

Packing:

Ostorin Tablet: Each box contains 30, 60 and 100 capsules in alu-alu blister pack.

Manufactured By:
The IBN SINA Pharmaceutical Industry PLC.
Shafipur, Gazipur, Bangladesh.