

FUNGIN

CLOTRIMAZOLE BP

Compositions:

Clotrimazole BP 1 % w/w.

Pharmacology:

Fungin Cream contains Clotrimazole BP, a synthetic antifungal agent, It is a broad spectrum antifungal agent and is used for the treatment of dermal infections caused by various species of pathogenic dermatophytes, yeasts and *Malassezia furfur*. The primary action of clotrimazole is against cell dividing and growing organisms. Absorption of clotrimazole in the skin is less than 0.5%.

Dosage And Administration:

Adequate Fungin Cream should be thinly applied 2-3 times daily to the diseased sites and rubbed in. After the prevention of the infection, it should be continued for 14 days.

Contraindications:

Clotrimazole is contraindicated in individuals who have shown hypersensitivity to any of its components.

Warning And Precaution:

If irritation or hypersensitivity develops with the use of clotrimazole, treatment should be discontinued and appropriate therapy should be instituted. The course of the cream should be completed although the symptoms are removed.

Side Effects:

A Small fraction of recipients, clotrimazole on the skin may causes stinging, erythema, edema, vesication, desquamation, pruritus and urticaria.

Use in Pregnancy and Lactation:

Infact a slight amount of medicine is absorbed in the blood when apply to the skin. Adequate examination had not done about the administration of the cream.It is unknown that wheather it is secreted to human milk and caution should be exercised when treating nursing mothers.

Drug Interaction:

No information is available.

Overdosage:

An overdose of clotrimazole topical is not expected to be dangerous. Seek emergency medical attention if anyone has accidentally swallowed the medication.

Storage:

Keep below 25°C temperature, away from light & moisture. Keep out of the reach of children.

Packing:

Fungin Cream : Tube containing 10 gm cream.

Manufactured By:

The IBN SINA Pharmaceutical Industry PLC.

Shafipur, Gazipur, Bangladesh.